

FLAT SEMILATTICES

G. GRÄTZER AND F. WEHRUNG

ABSTRACT. Let A , B , and S be $\{\vee, 0\}$ -semilattices and let $f: A \hookrightarrow B$ be a $\{\vee, 0\}$ -semilattice embedding. Then the canonical map, $f \otimes \text{id}_S$, of the tensor product $A \otimes S$ into the tensor product $B \otimes S$ is not necessarily an embedding.

The $\{\vee, 0\}$ -semilattice S is *flat*, if for every embedding $f: A \hookrightarrow B$, the canonical map $f \otimes \text{id}$ is an embedding. We prove that a $\{\vee, 0\}$ -semilattice S is flat if and only if it is distributive.

INTRODUCTION

Let A and B be $\{\vee, 0\}$ -semilattices. We denote by $A \otimes B$ the *tensor product* of A and B , defined as the free $\{\vee, 0\}$ -semilattice generated by the set

$$(A - \{0\}) \times (B - \{0\})$$

subject to the relations

$$\langle a, b_0 \rangle \vee \langle a, b_1 \rangle = \langle a, b_0 \vee b_1 \rangle,$$

for $a \in A - \{0\}$, $b_0, b_1 \in B - \{0\}$; and symmetrically,

$$\langle a_0, b \rangle \vee \langle a_1, b \rangle = \langle a_0 \vee a_1, b \rangle,$$

for $a_0, a_1 \in A - \{0\}$, $b \in B - \{0\}$.

$A \otimes B$ is a universal object with respect to a natural notion of *bimorphism*, see [2], [4], and [6]. This definition is similar to the classical definition of the tensor product of modules over a commutative ring. Thus, for instance, *flatness* is defined similarly: The $\{\vee, 0\}$ -semilattice S is *flat*, if for every embedding $f: A \hookrightarrow B$, the canonical map $f \otimes \text{id}_S: A \otimes S \rightarrow B \otimes S$ is an embedding.

Our main result is the following:

Theorem. *Let S be a $\{\vee, 0\}$ -semilattice. Then S is flat iff S is distributive.*

1. BACKGROUND

1.1. Basic concepts. We shall adopt the notation and terminology of [6]. In particular, for every $\{\vee, 0\}$ -semilattice A , we use the notation $A^- = A - \{0\}$. Note that A^- is a subsemilattice of A .

A semilattice S is *distributive*, if whenever $a \leq b_0 \vee b_1$ in S , then there exist $a_0 \leq b_0$ and $a_1 \leq b_1$ such that $a = a_0 \vee a_1$; equivalently, iff the lattice $\text{Id } S$ of all ideals of S , ordered under inclusion, is a distributive lattice; see [5].

Date: June 1, 1998.

1991 Mathematics Subject Classification. Primary 06B05, 06B10, 06A12, 08B25.

Key words and phrases. Tensor product, semilattice, lattice, antitone, flat.

The research of the first author was partially supported by the NSERC of Canada.

1.2. The set representation. In [6], we used the following representation of the tensor product.

First, we introduce the notation:

$$\nabla_{A,B} = (A \times \{0\}) \cup (\{0\} \times B).$$

Second, we introduce a partial binary operation on $A \times B$: let $\langle a_0, b_0 \rangle, \langle a_1, b_1 \rangle \in A \times B$; the *lateral join* of $\langle a_0, b_0 \rangle$ and $\langle a_1, b_1 \rangle$ is defined if $a_0 = a_1$ or $b_0 = b_1$, in which case, it is the join, $\langle a_0 \vee a_1, b_0 \vee b_1 \rangle$.

Third, we define bi-ideals: a nonempty subset I of $A \times B$ is a *bi-ideal* of $A \times B$, if it satisfies the following conditions:

- (i) I is hereditary;
- (ii) I contains $\nabla_{A,B}$;
- (iii) I is closed under lateral joins.

The *extended tensor product* of A and B , denoted by $A \overline{\otimes} B$, is the lattice of all bi-ideals of $A \times B$.

It is easy to see that $A \overline{\otimes} B$ is an algebraic lattice. For $a \in A$ and $b \in B$, we define $a \otimes b \in A \overline{\otimes} B$ by

$$a \otimes b = \nabla_{A,B} \cup \{ \langle x, y \rangle \in A \times B \mid \langle x, y \rangle \leq \langle a, b \rangle \}$$

and call $a \otimes b$ a *pure tensor*. A pure tensor is a principal (that is, one-generated) bi-ideal.

Now we can state the representation:

Proposition 1.1. *The tensor product $A \otimes B$ can be represented as the $\{\vee, 0\}$ -sub-semilattice of compact elements of $A \overline{\otimes} B$.*

1.3. The construction of $A \overline{\otimes} B$. The proof of the Theorem uses the following representation of the tensor product, see J. Anderson and N. Kimura [1].

Let A and B be $\{\vee, 0\}$ -semilattices. Define

$$A \vec{\otimes} B = \text{Hom}(\langle A^-; \vee \rangle, \langle \text{Id } B; \cap \rangle)$$

and for $\xi \in A \vec{\otimes} B$, let

$$\varepsilon(\xi) = \{ \langle a, b \rangle \in A^- \times B^- \mid b \in \xi(a) \} \cup \nabla_{A,B}.$$

Proposition 1.2. *The map ε is an order preserving isomorphism between $A \overline{\otimes} B$ and $A \vec{\otimes} B$ and, for $H \in A \overline{\otimes} B$, $\varepsilon^{-1}(H)$ is given by the formula*

$$\varepsilon^{-1}(H)(a) = \{ b \in B \mid \langle a, b \rangle \in H \},$$

for $a \in A^-$.

If $a \in A$ and $b \in B$, then $\varepsilon(a \otimes b)$ is the map $\xi: A^- \rightarrow \text{Id } B$:

$$\xi(x) = \begin{cases} (b], & \text{if } x \leq a; \\ \{0\}, & \text{otherwise.} \end{cases}$$

If A is *finite*, then a homomorphism from $\langle A^-; \vee \rangle$ to $\langle \text{Id } B; \cap \rangle$ is determined by its restriction to $J(A)$, the set of all join-irreducible elements of A . For example, let A be a finite Boolean semilattice, say $A = P(n)$ (n is a non-negative integer, $n = \{0, 1, \dots, n-1\}$), then $A \overline{\otimes} B \cong (\text{Id } B)^n$, and the isomorphism from $A \overline{\otimes} B$ onto $(\text{Id } B)^n$ given by Proposition 1.2 is the unique complete $\{\vee, 0\}$ -homomorphism sending every element of the form $\{i\} \otimes b$ ($i < n$ and $b \in B$) to $\langle (\delta_{ij} b] \mid j < n \rangle$

(where δ_{ij} is the Kronecker symbol). If $n = 3$, let $\beta: P(3) \otimes S \rightarrow (\text{Id } S)^3$ denote the natural isomorphism.

Next we compute $A \otimes B$, for $A = M_3$, the diamond, and $A = N_5$, the pentagon (see Figure 1). In the following two subsections, let S be a $\{\vee, 0\}$ -semilattice. Furthermore, we shall denote by $\tilde{S}$ the ideal lattice of S , and identify every element s of S with its image, (s) , in $\tilde{S}$.




Figure 1

1.4. The lattices $M_3 \otimes S$ and $M_3[\tilde{S}]$; the map i . Let $M_3 = \{0, p, q, r, 1\}$, $J(M_3) = \{p, q, r\}$ (see Figure 1). The nontrivial relations of $J(M_3)$ are the following:

$$p < q \vee r, \quad q < p \vee r, \quad \text{and} \quad r < p \vee q. \quad (1)$$

Accordingly, for every lattice L , we define

$$M_3[L] = \{ \langle x, y, z \rangle \in L^3 \mid x \wedge y = x \wedge z = y \wedge z \} \quad (2)$$

(this is the *Schmidt's construction*, see [9] and [10]). The isomorphism from $M_3 \otimes S$ onto $M_3[\tilde{S}]$ given by Proposition 1.2 is the unique complete $\{\vee, 0\}$ -homomorphism α such that, for all $x \in S$,

$$\begin{aligned} \alpha(p \otimes x) &= \langle x, 0, 0 \rangle, \\ \alpha(q \otimes x) &= \langle 0, x, 0 \rangle, \\ \alpha(r \otimes x) &= \langle 0, 0, x \rangle. \end{aligned}$$

We shall make use later of the unique $\{\vee, 0\}$ -embedding

$$i: M_3 \hookrightarrow P(3)$$

defined by

$$\begin{aligned} i(p) &= \{1, 2\}, \\ i(q) &= \{0, 2\}, \\ i(r) &= \{0, 1\}. \end{aligned}$$

1.5. The lattices $N_5 \overline{\otimes} S$ and $N_5[\tilde{S}]$; the map i' . Let $N_5 = \{0, a, b, c, 1\}$, $J(N_5) = \{a, b, c\}$ with $a > c$ (see Figure 1). The nontrivial relations of $J(N_5)$ are the following:

$$c < a \quad \text{and} \quad a < b \vee c. \quad (3)$$

Accordingly, for every lattice L , we define

$$N_5[L] = \{ \langle x, y, z \rangle \in L^3 \mid y \wedge z \leq x \leq z \}. \quad (4)$$

The isomorphism from $N_5 \overline{\otimes} S$ onto $N_5[\tilde{S}]$, given by Proposition 1.2, is the unique complete $\{\vee, 0\}$ -homomorphism α' such that, for all $x \in S$,

$$\begin{aligned} \alpha'(a \otimes x) &= \langle x, 0, x \rangle, \\ \alpha'(b \otimes x) &= \langle 0, x, 0 \rangle, \\ \alpha'(c \otimes x) &= \langle 0, 0, x \rangle. \end{aligned}$$

We shall make use later of the unique $\{\vee, 0\}$ -embedding

$$i': N_5 \hookrightarrow P(3)$$

defined by

$$\begin{aligned} i'(a) &= \{0, 2\}, \\ i'(b) &= \{1, 2\}, \\ i'(c) &= \{0\}. \end{aligned}$$

1.6. The complete homomorphisms $f \overline{\otimes} g$. The proof of the following lemma is straightforward:

Lemma 1.3. *Let A , B , A' , and B' be $\{\vee, 0\}$ -semilattices, let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be $\{\vee, 0\}$ -homomorphisms. Then the natural $\{\vee, 0\}$ -homomorphism $h = f \otimes g$ from $A \otimes B$ to $A' \otimes B'$ extends to a unique complete $\{\vee, 0\}$ -homomorphism $\bar{h} = f \overline{\otimes} g$ from $A \overline{\otimes} B$ to $A' \overline{\otimes} B'$. Furthermore, if h is an embedding, then $\bar{h}$ is also an embedding.*

We refer to Proposition 3.4 of [6] for an explicit description of the map $\bar{h}$.

2. CHARACTERIZATION OF FLAT $\{\vee, 0\}$ -SEMI LATTICES

Our definition of flatness is similar to the usual one for modules over a commutative ring:

Definition. A $\{\vee, 0\}$ -semilattice S is *flat*, if for every embedding $f: A \hookrightarrow B$ of $\{\vee, 0\}$ -semilattices, the tensor map $f \otimes \text{id}_S: A \otimes S \rightarrow B \otimes S$ is an embedding.

In this definition, id_S is the identity map on S .

In Lemmas 2.1–2.3, let S be a $\{\vee, 0\}$ -semilattice and we assume that both homomorphisms $f = i \otimes \text{id}_S$ and $f' = i' \otimes \text{id}_S$ are embeddings.

As in the previous section, we use the notation $\tilde{S} = \text{Id } S$, and identify every element s of S with the corresponding principal ideal $[s]$.

We define the maps $g: M_3[\tilde{S}] \rightarrow \tilde{S}^3$ and $g': N_5[\tilde{S}] \rightarrow \tilde{S}^3$ by the following formulas:

$$\begin{aligned} \text{For all } \langle x, y, z \rangle \in M_3[\tilde{S}], \quad & g(\langle x, y, z \rangle) = \langle y \vee z, x \vee z, x \vee y \rangle, \\ \text{For all } \langle x, y, z \rangle \in N_5[\tilde{S}], \quad & g'(\langle x, y, z \rangle) = \langle z, y, x \vee y \rangle. \end{aligned}$$

Note that g and g' are complete $\{\vee, 0\}$ -homomorphisms. The proof of the following lemma is a straightforward calculation.

Lemma 2.1. *The following two diagrams commute:*

$$\begin{array}{ccc}
 M_3 \otimes S & \xrightarrow{f} & P(3) \otimes S \\
 \alpha \downarrow & & \downarrow \beta \\
 M_3[\tilde{S}] & \xrightarrow{g} & \tilde{S}^3 \\
 \\
 N_5 \otimes S & \xrightarrow{f'} & P(3) \otimes S \\
 \alpha' \downarrow & & \downarrow \beta \\
 N_5[\tilde{S}] & \xrightarrow{g'} & \tilde{S}^3
 \end{array}$$

Therefore, both g and g' are embeddings.

Lemma 2.2. *The lattice $\tilde{S}$ does not contain a copy of M_3 .*

Proof. Suppose, on the contrary, that $\tilde{S}$ contains a copy of M_3 , say $\{o, x, y, z, i\}$ with $o < x, y, z < i$. Then both elements $u = \langle x, y, z \rangle$ and $v = \langle i, i, i \rangle$ of L^3 belong to $M_3[\tilde{S}]$, and $g(u) = g(v) = \langle i, i, i \rangle$. This contradicts the fact, proved in Lemma 2.1, that g is one-to-one. $\square$

Lemma 2.3. *The lattice $\tilde{S}$ does not contain a copy of N_5 .*

Proof. Suppose, on the contrary, that $\tilde{S}$ contains a copy of N_5 , say $\{o, x, y, z, i\}$ with $o < x < z < i$ and $o < y < i$. Then both elements $u = \langle x, y, z \rangle$ and $v = \langle z, y, z \rangle$ of L^3 belong to $N_5[\tilde{S}]$, and $g'(u) = g'(v) = \langle z, y, i \rangle$. This contradicts the fact, proved in Lemma 2.1, that g' is one-to-one. $\square$

Lemmas 2.2 and 2.3 together prove that $\tilde{S}$ is distributive, and therefore S is a distributive semilattice. Now we are in position to prove the main result of this paper in the following form:

Theorem 1. *Let S be a $\{\vee, 0\}$ -semilattice. Then the following are equivalent:*

- (i) S is flat.
- (ii) Both homomorphisms $i \otimes \text{id}_S$ and $i' \otimes \text{id}_S$ are embeddings.
- (iii) S is distributive.

Proof.

(i) *implies* (ii). This is trivial.

(ii) *implies* (iii). This was proved in Lemmas 2.2 and 2.3.

(iii) *implies* (i). Let S be a distributive $\{\vee, 0\}$ -semilattice; we prove that S is flat. Since the tensor product by a fixed factor preserves direct limits (see Proposition 2.6 of [6]), flatness is preserved under direct limits. By P. Pudlák [8], every distributive join-semilattice is the direct union of all its finite distributive subsemilattices; therefore, it suffices to prove that every finite distributive $\{\vee, 0\}$ -semilattice S is flat. Since S is a distributive lattice, it admits a lattice embedding into a finite Boolean lattice B . We have seen in Section 1.3 that if $B = P(n)$, then $A \otimes B = A^n$ (up to a natural isomorphism), for every $\{\vee, 0\}$ -semilattice A . It follows that B

is flat. Furthermore, the inclusion map $S \hookrightarrow B$ is a lattice embedding; in particular, with the terminology of [6], an *L-homomorphism*. Thus, the natural map from $A \otimes S$ to $A \otimes B$ is, by Proposition 3.4 of [6], a $\{\vee, 0\}$ -semilattice embedding. This implies the flatness of S . $\square$

3. DISCUSSION

It is well-known that a module over a given principal ideal domain R is flat if and only if it is torsion-free, which is equivalent to the module being a direct limit of (finitely generated) free modules over R . So the analogue of the concept of torsion-free module for semilattices is the concept of distributive semilattice. This analogy can be pushed further, by using the following result, proved in [3]: *a join-semilattice is distributive iff it is a direct limit of finite Boolean semilattices*.

Problem 1. Let $\mathbf{V}$ be a variety of lattices. Let us say that a $\{\vee, 0\}$ -semilattice S is *in* $\mathbf{V}$, if $\text{Id } S$ as a lattice is in $\mathbf{V}$. Is every $\{\vee, 0\}$ -semilattice in $\mathbf{V}$ a direct limit (resp., direct union) of *finite* join-semilattices in $\mathbf{V}$?

If $\mathbf{V}$ is the variety of all lattices, we obtain the obvious result that every $\{\vee, 0\}$ -semilattice is the direct union of its finite $\{\vee, 0\}$ -subsemilattices. If $\mathbf{V}$ is the variety of all distributive lattices, there are two results (both quoted above): P. Pudlák's result and K. R. Goodearl and the second author's result.

Problem 2. Let $\mathbf{V}$ be a variety of lattices. When is a $\{\vee, 0\}$ -semilattice S flat with respect to $\{\vee, 0\}$ -semilattice embeddings in $\mathbf{V}$? That is, when is it the case that for all $\{\vee, 0\}$ -semilattices A and B in $\mathbf{V}$ and every semilattice embedding $f: A \hookrightarrow B$, the natural map $f \otimes \text{id}_S$ is an embedding?

REFERENCES

- [1] J. Anderson and N. Kimura, *The tensor product of semilattices*, Semigroup Forum **16** (1978), 83–88.
- [2] G. Fraser, *The tensor product of semilattices*, Algebra Universalis **8** (1978), 1–3.
- [3] K. R. Goodearl and F. Wehrung, *Representations of distributive semilattices by dimension groups, regular rings, C^* -algebras, and complemented modular lattices*, submitted for publication, 1997.
- [4] G. Grätzer, H. Lakser, and R. W. Quackenbush, *The structure of tensor products of semilattices with zero*, Trans. Amer. Math. Soc. **267** (1981), 503–515.
- [5] G. Grätzer, *General Lattice Theory. Second Edition*, Birkhäuser Verlag, Basel, 1998. xix+663 pp.
- [6] G. Grätzer and F. Wehrung, *Tensor products of semilattices with zero, revisited*, submitted for publication, 1998.
- [7] ———, *Tensor products and transferability of semilattices*, submitted for publication, 1998.
- [8] P. Pudlák, *On congruence lattices of lattices*, Algebra Universalis **20** (1985), 96–114.
- [9] R. W. Quackenbush, *Non-modular varieties of semimodular lattices with a spanning M_3* . Special volume on ordered sets and their applications (L'Arbresle, 1982). Discrete Math. **53** (1985), 193–205.
- [10] E. T. Schmidt, *Zur Charakterisierung der Kongruenzverbände der Verbände*, Mat. Časopis Sloven. Akad. Vied **18** (1968), 3–20.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, MANITOBA, CANADA
R3T 2N2

E-mail address: `gratzer@cc.umanitoba.ca`

URL: `http://www.maths.umanitoba.ca/homepages/gratzer.html/`

C.N.R.S., DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE CAEN, 14032 CAEN CEDEX,
FRANCE

E-mail address: `gremlin@math.unicaen.fr`

URL: `http://www.math.unicaen.fr/~wehrung`